

UCRL-93519
PREPRINT

BETA-DELAYED FISSION CALCULATIONS
FOR THE ASTROPHYSICAL r - PROCESS

B.S. Meyer
W.M. Howard
G.J. Mathews
P. Moller
K. Takahashi

This paper was prepared for submittal to
ACS Symposium on Recent Advances in the
Study of Nuclei off the Line of Stability
Chicago, Illinois
September 8-13, 1985

October 9, 1985

The logo of the Lawrence Livermore National Laboratory is a large, stylized 'V' shape. The 'V' is composed of several horizontal layers of different patterns: a white top layer, a dark grey middle layer, and a black bottom layer. The text 'Lawrence Livermore National Laboratory' is written diagonally across the right side of the 'V' in a sans-serif font.

Lawrence
Livermore
National
Laboratory

This is a preprint of a paper intended for publication in a journal or proceedings. Since changes may be made before publication, this preprint is made available with the understanding that it will not be cited or reproduced without the permission of the author.

DISCLAIMER

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial products, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

BETA-DELAYED FISSION CALCULATIONS FOR THE ASTROPHYSICAL r-PROCESS

B.S.Meyer, W.M.Howard, G.J.Mathews, P.Möller and K.Takahashi

University of California, Lawrence Livermore National Laboratory, Livermore, CA 94550

Abstract

We discuss RPA calculations of the Gamow-Teller properties of neutron-rich nuclei to study the effect of β -delayed fission and neutron emission on the production of Th, U and Pu chronometric nuclei in the astrophysical r-process. We find significant differences in the amount of β -delayed fission when compared with the recent calculations of Thielemann et al. (1983). In the simplest case of a constant abundance along the r-process path, however, the inferred production ratios in both calculations are similar.

The study of the decay of nuclei from the r-process path back to the β -stability line is an area of astrophysics which requires knowledge of β -strength functions off the line of beta stability. This knowledge is especially important for the determination of the abundances of the progenitors of the Th-U-Pu chronometers since β -delayed fission and neutron emission during decay back to the stability line may significantly affect the final abundance distribution of these nuclei ([BER69], [WEN75], [KOD75], [KRUS1]). The β -strength function for nuclei along the decay back paths [coupled with neutron separation energies (S_n), fission barrier heights (B_f) and β -decay Q-values (Q_β)] determines the amount of β -delayed fission and neutron emission that occurs during the cascade back to the β -stability line.

A recent analysis by Thielemann et al. [THI83] of the effects of β -delayed processes on the progenitors of the Th-U-Pu chronometers showed that these processes (delayed fission in particular) did indeed significantly influence the final abundances of the chronometer progenitors. This leads to a long age for the Galaxy. In view of the importance of this result, it is useful to re-examine the calculation with a nuclear model that includes the effects of nuclear deformation on the β -decay rates, fission barriers, and neutron separation energies self-consistently.

Since many of the nuclei involved are presumably highly-deformed, we have used the Nilsson RPA code of Krumlinde and Möller [KRUS84], which calculates β -strength functions, using an infinite-range residual (Gamow-Teller) interaction with a strength $\chi_{GT} = 23/A$ MeV. With this code, we have calculated β -strength functions for 118 nuclei lying in the mass range from $A=232$ to $A=255$ and from the line of β -stability to the r-process path given by Thielemann et al. [THI83] (see Fig.1). These 118 nuclei satisfied the criteria for possible β -delayed fission (Q_β of precursor $\geq B_f$ of emitter) and/or possible β -delayed neutron emission (Q_β of precursor $\geq S_n$ of emitter) as determined from the values given in [HOW80].

The required input values are the deformation parameters ϵ_2 and ϵ_4 , and the numbers κ and μ which determine the size of the $l \cdot s$ force and l^2 terms for the harmonic oscillator. We adopted the values $\kappa_p = 0.0577$ and $\mu_p = 0.650$ for protons and $\kappa_n = 0.0635$ and $\mu_n = 0.325$ for neutrons over the entire range studied, in accordance with [HOW80]. The values of ϵ_2 and ϵ_4 for each nucleus were taken from the same source.

From our β -strength distributions, we then calculated rates as a function of excitation energy in the daughter nucleus. These rates and the Q_β , B_f , and S_n values [HOW80] give the amount of β -delayed fission and neutron emission for each daughter nucleus. Our results are shown in Fig. 1. The numbers in the squares in Fig. 1(a) give the percentage of decays resulting in a daughter nucleus excitation energy greater than or equal to the fission barrier height. The numbers in the squares in Fig. 1(b) give the percentage of decays resulting in an daughter nucleus excitation energy greater than or equal to the neutron separation energy but less than the fission barrier height. In order to determine the maximum possible effect of delayed fission, we take the number from (a) as the amount of β -delayed fission and the number from (b) as the amount of β -delayed neutron emission in the daughter nucleus. This undoubtedly leads to an overestimate of delayed fission probabilities in some cases.

For the purpose of simple comparison with [THI83], we assume constant abundances (1.0 per isobar) along the r-process path shown in Fig.1. After β -delayed fission and neutron emission during decay back, the final abundances are those shown in Table 1. Clearly β -delayed fission plays a large role in decay back at $A \leq 250$. On the other hand, β -delayed neutron emission does not affect the abundances too much, since loss from chain A is more or less compensated for by gain from chain A+1, except in a few cases ($A = 234, 236, 239, 244, 248$, and 249). Of course, a more significant effect of delayed neutron emission will be seen if we take more realistic initial abundances along the r-process path, which usually exhibit a strong even-odd effect.




Fig. 1(a). Calculated maximum possible values of β -delayed fission probabilities (in %) shown for the precursor nuclei. The arrow follows the r-process path given in [THI83]. The crosses indicate β -stable nuclei, and the dashed lines indicate α decays. The r-process nuclear cosmo-chronometers are circled.




Fig. 1(b). Calculated minimum possible values of β -delayed neutron emission probabilities (in %). See the caption to Fig.1(a).

Table 1. Isobaric abundance changes during the cascade from the initial value of 100 % at the r-process path shown in Fig. 1 down to the β -stable nuclei marked by crosses.

A	%	A	%
232	78	244	47
233	102	245	112
234	70	246	110
235	118	247	124
236	133	248	64
237	92	249	64
238	97	250	47
239	67	251	32
240	115	252	67
241	111	253	18
242	103	254	11
243	94	255	6

From the results in Table 1, we can find the r-process production ratios of interest to nuclear cosmo-chronology: $^{232}\text{Th}/^{238}\text{U}$, $^{235}\text{U}/^{238}\text{U}$, and $^{244}\text{Pu}/^{238}\text{U}$. These are obtained by summing the α -decay progenitors for each isotope and considering the leak due to spontaneous fission. Our results are $^{232}\text{Th}/^{238}\text{U} = 1.60$, $^{235}\text{U}/^{238}\text{U} = 1.42$, $^{244}\text{Pu}/^{238}\text{U} = 0.57$. Our value for $^{232}\text{Th}/^{238}\text{U}$ ratio in this approximation of constant r-process abundances would imply a lower limit on the Galaxy's age of 8.8 Gyr. However, it is important to note that the final production ratios will be dependent on the explicit r-

process calculation employed [FOW85]. (See also [MEY85], [THI83] and [YOK83] for further details on nuclear cosmo-chronology.)

For comparison, we give the values for the chronometer production ratios determined by Thielemann et al. [THI83] for constant abundances along the r-process path: $^{232}\text{Th}/^{238}\text{U} = 1.63$, $^{235}\text{U}/^{238}\text{U} = 1.21$, and $^{244}\text{Pu}/^{238}\text{U} = 0.13$. To our surprise, our $^{232}\text{Th}/^{238}\text{U}$ ratio agrees well with [THI83], despite the large difference between the two calculations in the amount of β -delayed fission (compare Fig. 1(a) with Fig. 2 of [THI83]). This result suggests that β -delayed processes do indeed significantly affect the abundances of the progenitors of the r-process chronometers. The other two sets of ratios differ rather strongly, however. These differences are not surprising because of the different β -strength functions and adopted mass formulae (for Q_β and S_n), although both sets of calculations used the same fission barrier heights. It is worth emphasizing that the virtue of our calculations are that all input data were taken from a single source [HOW80], i.e., the effects of nuclear deformations on the fission barriers and β -strength functions were treated self-consistently. We understand that many uncertainties still exist in this calculation and, therefore, that further work is still required. We hope to pursue this study in the near future and will be aided by improvements in the code used for calculating the β -strength functions (e.g., the inclusion of the first-forbidden decay) and in the set of Q_β , B_f and S_n values. We will also include a more realistic determination of the competition between delayed fission and delayed neutron emission. These improvements should lead to a better understanding of decay back to the β -stability line and help in the search for the astrophysical site(s) for the r-process.

We express our thanks to Prof. William A. Fowler for useful discussions and encouragement.

Work performed under auspices of the U.S. Department of Energy by the Lawrence Livermore National Laboratory under contract number W-7405-ENG-48, and supported in part by the LLNL Institute for Geophysics and Planetary Science. One of us (BSM) acknowledges the support of National Science Foundation Graduate Fellowship.

References

- [BER69] E.Yu.Berlovich and Yu.N.Novikov, Phys.Lett. 29B 155 (1969)
- [FOW85] W.A.Fowler and C.C.Meisel, in Proc.Symp. on Cosmogonical Processes, Boulder, Colorado, 1985, in press
- [HOW80] W.M.Howard and P.Möller, Atm.Nucl.Dat.Tables 25 219 (1980)
- [KOD75] T.Kodama and K.Takahashi, Nucl.Phys. A239 489 (1975)
- [KRU81] J.Krumlinde, P.Möller, C.-O.Wene and W.M.Howard, in Proc.4th Int.Conf. on NUclei Far From Stability, Helsingör, 1981; CERN Rpt. 81-09, p.260
- [KRU84] J.Krumlinde and P.Möller, Nucl.Phys. A417 419 (1984)
- [MEY85] B.S.Meyer and D.N.Schramm, in Proc.5th Moriond Astrophysics Meeting on Nucleosynthesis and its Implications on Nuclear and Particle Physics, Les Arcs, 1985 (Reidel) in press
- [THI83] F.-K.Thielemann, J.Metzinger and H.V.Klapdor, Astron.Astrophys. 123 162 (1983)
- [WEN75] C.-O.Wene, Astron.Astrophys. 44 233 (1975)
- [YOK83] K.Yokoi, K.Takahashi and M.Arnoold, Astron.Astrophys. 118 6 (1983)